



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

In the triangles FCB and FBD , $\angle FCB = \angle FBA$, since arc $AF = \text{arc } FB$; also $\angle CFB$ is common, hence the triangles are similar, and $FC : FB = FB : FD$; but $FL (= FC) : FB = FB : FH$. Therefore $FH = FD$ and $HL = CD$.

Hence in the triangle ABC , AB is the given base, $\angle ACB$ the given vertical angle, and CD the given bisector, and the triangle is satisfied in every condition.

Also solved by L. E. Newcomb, and A. H. Holmes.

CALCULUS.

233. Proposed by W. J. GREENSTREET, M. A., Editor of the Mathematical Gazette, Stroud, England.

Prove that
$$\int_0^\infty \frac{x^{a-1} dx}{1+2x \cos \theta + x^2} = \frac{\pi \sin(1-a)\theta}{\sin a\pi \sin \theta}.$$

Solution by G. B. M. ZERR, A. M., Ph. D., Parsons, W. Va.

$$\begin{aligned} \int_0^\infty \frac{x^{a-1} dx}{1+2x \cos \theta + x^2} &= \frac{1}{a} \int_0^\infty \frac{dx}{\sin^2 \theta + (x + \cos \theta)^2} \\ &= \frac{1}{a \sin \theta} \tan^{-1} \left(\frac{x + \cos \theta}{\sin \theta} \right) \Big|_0^\infty = \frac{\theta}{a \sin \theta} \end{aligned}$$

The problem giving the result stated is as follows:

$$\begin{aligned} \int_0^\infty \frac{x^{a-1} dx}{1+2x \cos \theta + x^2} &= \frac{1 - \sqrt{(-1) \cot \theta}}{2} \int_0^\infty \frac{x^{a-2} dx}{x + \cos \theta + \sqrt{(-1) \sin \theta}} \\ &\quad + \frac{1 + \sqrt{(-1) \cot \theta}}{2} \int_0^\infty \frac{x^{a-2} dx}{x + \cos \theta - \sqrt{(-1) \sin \theta}} = P. \end{aligned}$$

Let $x = y[\cos \theta \pm \sqrt{(-1) \sin \theta}]$, the plus sign for the first term, the minus sign for the second term.

$$\begin{aligned} \therefore P &= \frac{1}{2} [1 - \sqrt{(-1) \cot \theta}] [\cos(a-2)\theta + \sqrt{(-1) \sin(a-2)\theta}] \int_0^\infty \frac{y^{a-2} dy}{1+y} \\ &\quad + \frac{1}{2} [1 + \sqrt{(-1) \cot \theta}] [\cos(a-2)\theta - \sqrt{(-1) \sin(a-2)\theta}] \int_0^\infty \frac{y^{a-2} dy}{1+y}. \\ \therefore P &= [\cos(a-2)\theta + \sin(a-2)\theta \cot \theta] \int_0^\infty \frac{y^{a-2} dy}{1+y} \\ &= \frac{\sin(a-1)\theta}{\sin \theta} \int_0^\infty \frac{y^{a-2} dy}{1+y} = \frac{\sin(a-1)\theta}{\sin \theta} \cdot \frac{\pi}{\sin(a-1)\pi}. \\ \therefore P &= \frac{\pi \sin(1-a)\theta}{\sin \theta \sin a\pi} \end{aligned}$$

This problem was incorrectly stated, the error being due to an oversight in reading proof. It is correctly stated above, the numerator being x^{a-1} instead of a^{-1} . ED. F.